

51st Scientific and Technical Subcommittee – UN COPUOS
Vienna , 12 February 2014


HIGHLIGHTS OF THE SCOSTEP CAUSES II SCIENTIFIC PROGRAM (2009-2013) - UNDERSTANDING THE CLIMATE AND WEATHER OF THE SUN-EARTH SYSTEM

Marianna G. Shepherd
Scientific Secretary

Scientific Committee on Solar-Terrestrial Physics (SCOSTEP)



HISTORY OF SCOSTEP SCIENCE PROGRAMS




CAWSES: CLIMATE AND WEATHER OF THE SUN-EARTH SYSTEM

- CAWSES was **established in 2004** as SCOSTEP's international program to link the world's scientists in a cooperative effort in improving our understanding of the Solar Terrestrial relations and their impact on life and society.
- **Special emphasis** was given on the **short- (weather) and long-term (climate) variabilities** of the solar activity and their effects/responses in Geospace and Earth's environment.
- CAWSES combined both **purely scientific and application motivations** for enhancing our understanding of variations in the Sun-Earth system.
- Ground-based and satellite observations, and numerical modeling.
- **Capacity building activities** involved researchers in developing countries, together with educational opportunities for students.


CAWSES

Phase 1: 2004-2008



CAWSES II

Phase 2: 2009-2013




CAWSES II 2009-2013

- Fundamental questions of how the coupled Sun-Earth system operates on timescales of minutes to millennia
- Questions that require coordinated inter-disciplinary international effort

Four major tasks of CAWSES II:

- What is the solar influence on the Earth's climate?
- How will geospace respond to an altered climate?
- How does short-term solar variability affect the geospace environment?
- What is the geospace response to variable inputs from the lower atmosphere?



CAWSES CAWSES II

(during the rising phase of Cycle 24)

CAWSES II Task Groups

Solar influences on the Earth's climate

TG1

TG 3: Solar variability effects on geospace environment

TG2



Effects of climate on geospace

TG4

Variable input from lower atmosphere on geospace

Capacity building

Informatics and eScience





WHAT ARE THE SOLAR INFLUENCES ON THE EARTH'S CLIMATE?

- **Solar variability drives the Earth's environment** on time scales ranging from minutes to millennia. **Feedbacks are inherent in the Earth system** and may amplify the direct forcing effects from the Sun.
- **The influence of this solar variability on Earth's climate** is a key issue of IPCC, and one that continues to be highlighted by policy makers, climate change skeptics, and the media.

OPEN QUESTIONS

- What is the importance of **spectral variations** to solar influences on climate?
- What is the effect of **energetic particle** forcing on the whole atmosphere and what are the implications for climate?
- How well do **models reproduce and predict solar irradiance and energetic particle influences** on the atmosphere and climate?

SCIENTIFIC PROGRESS

■ Importance of spectral variations to solar influences on climate

- Solar influence on climate - an important contribution to climate variability → Shift focus from global to regional response
- Importance of top-down stratospheric UV mechanism accepted
- Improved *Total Solar Irradiance* and *Solar Spectral Irradiance (SSI)* measurements: solar cycle variation estimates → New value of the "solar constant" (1361 Wm^{-2})
- Direct effects of SSI – Ionization rates; Chemical changes: NO_y , O_3 , some other constituents; Radiative effects

■ The effect of energetic particle forcing on the whole atmosphere and its implications for climate


- Strong indirect effect of energetic particle precipitation (EPP) on stratosphere; studies of effects on troposphere and surface – inconclusive.
- Solar protons impact polar chemistry, but no long-term (>years) effect

■ Modeling of solar irradiance and energetic particle influences on the atmosphere and climate

- Global models start including galactic cosmic rays (GCR) ionization - CERN CLOUD measurements show GCR stimulate aerosol nucleation at upper tropospheric temperatures → Coincident surface-based measurements of aerosol nucleation and ion contribution




WHAT IS THE GEOSPACE RESPONSE TO VARIABLE WAVES FROM THE LOWER ATMOSPHERE?

- A variety of new evidence suggests that **tropospheric weather is an important ingredient in space weather**. Equatorial ionospheric densities are modulated by **atmospheric waves** driven by persistent **tropical rainstorms**. Radio waves generated by **lightning strokes** in rainstorms **interact with radiation belt particles** to clear a "safe" zone between the inner and outer belts in the magnetosphere.
- **Atmospheric gravity waves** generated by hurricanes and typhoons may seed **plasma bubbles** in the low latitude ionosphere → The extent to which the effects of this quiescent **atmospheric variability are transmitted to the magnetosphere** is yet to be resolved.
- **Overall GOAL:** Elucidate the **dynamical coupling** from the lower atmosphere to the geospace (upper atmosphere, ionosphere, and magnetosphere), for various frequencies and scales, such as **gravity waves, tides, and planetary waves, and for equatorial, middle and high latitudes.**



**COSMIC/
Formosat 3C**
Northern EIA region


**F-region
(250 - 450 km)**



**E-region
Dynamo Modulation**




**MLT/E-region
(80 - 140 km)**




T, u, v, ρ
Forbes et al., 2006;
Oberheide et al., 2005

“Wave-4”, DE3, SE2, ...
Non-migrating tides



Nitric Oxide (NOx)
Oberheide and Forbes, 2008

Tropical Troposphere





**weather
satellites**

Deep Convective Clouds
ISCCP

TEC Observations with GPS (GEONET; dense GPS network in Japan, 1300 stations)

Tsunami, occurred on March 11, 2011, generated **sound waves and gravity waves**, that propagated up to the ionosphere causing **TEC perturbations**.



WHAT WAS ACHIEVED?

- Collaborative effort by the neutral atmosphere and plasma communities leading to a much better understanding of how neutral dynamics originating in the lower atmosphere impact Earth's ionosphere.
- A significant progress in separating ionospheric variability introduced by the driving from below and from the Sun.
- The specific contributions were obtained from five observational campaigns and the support of dedicated workshops resulting in special issues in the peer-reviewed literature.
- **OVERALL OUTCOME : *Results show significant penetration of atmospheric waves into the ionosphere in all frequency ranges: acoustic waves, gravity waves, tides, and planetary waves***
- **Current Science Challenges (among others):**
 - Direct penetration of atmospheric waves into thermosphere and then into the ionosphere
 - Electromagnetic coupling from E (~100 km) to F (~300 km) region and then to thermosphere
- Impact of Low Solar Activity: *Enhanced wave effects, due to (i) cooler thermosphere and reduced wave dissipation, and (ii) less solar induced disturbances*

HOW WILL THE GEOSPACE RESPOND TO AN ALTERED CLIMATE?



- Radiative, chemical, and dynamical forcing from below contributes to disturbances of the upper atmosphere.
- Patterns of cooling and contraction of the upper atmosphere in response to rising greenhouse gas concentrations are emerging from model studies and observations, consistent with a strong connection to changes in the lower atmosphere.
- Rising greenhouse gas concentrations alter the ionosphere in a variety of ways and could be transmitted to the magnetosphere.
- These changes may have unforeseen consequences for space-related technologies and societal infrastructures.

SCIENTIFIC HIGHLIGHTS

- 33-year PMC (Polar Mesospheric Clouds) data (at ~83 km) from the SBUV satellite → significantly smaller trends than previously reported. Long-term increases in PMC brightness, particularly in the Northern Hemisphere.
- 50-year data of NLC (Noctilucent Clouds, observed from the ground) → a weak positive trend (but statistically insignificant) in their appearance frequency - results consistent with satellite observations and models → the trend's magnitude increases with latitude.
- Models show that CO₂, CH₄ and CFCs drive mesospheric trends. Since ~1997, no significant trends are found in mesospheric temperature and PMC frequency, apparently due to the stabilization of stratospheric ozone (Montreal protocol).
- Onset of the Antarctic PMC season - highly correlated to the time of breakdown of the stratospheric winter vortex (*intra-hemispheric coupling*).
- Seven documented cases of impulsive increases of PMC resulting from poleward transport of water vapor from the exhaust of the U.S. Space Shuttle main engine.

TRENDS IN TOTAL ELECTRON CONTENT (TEC)


TEC trends from global GPS observations over 1995-2010 (Lean et al., 2011).



Daily averaged global trend **+0.6 TECU/decade** (if solar EUV flux in 2008 is the same as in 1996) or **+3.2 TECU/decade** (if solar EUV flux in 2008 is 15% lower than in 1996). The latter trend is not physically plausible; the former may be questioned (according to models and foF2 trends the expected trend should be negative).

➔ **Impact of peculiar solar cycle 23?**

Thermospheric neutral density trends near 400km derived from satellite drag over three solar cycles – Emmert et al. (2008).



Trends are negative for any level of solar activity but they are sub-stancially stronger under solar minimum conditions due to larger relative role of CO₂ in radiative cooling.

OPEN QUESTIONS

- **Trend drivers** (GHGs, solar/geomagnetic, ozone, water vapor) **are changing** → these need to be understood/quantified.
- What are the **trends in the atmospheric dynamics** (circulation and wave activity)?
- **Improve accuracy in quantifying trends** in various parameters, and to reduce the model-observation trend differences.
- On longer-time scale – to **join upper atmospheric trends with long-term changes in the stratosphere** into one scenario.

IMPACT OF LONG-TERM CHANGES & TRENDS ON SATELLITE-BASED TECHNOLOGIES

- Longer orbital **life-time of satellites** and of **dangerous space debris** (more collisions with satellites) in response to cooling and contracting thermosphere.
- Long-term changes in **TEC and gravity wave activity** could have impact on ionospheric influence on GNSS (GPS, GALILEO, GLONASS etc.) signal propagation + ionospheric corrections and signal stability.

HOW DOES SHORT-TERM SOLAR VARIABILITY AFFECT THE GEOSPACE ENVIRONMENT?

■ CME-ICME connection

- **Advanced modeling** for solar eruption and CME release
- **Observed & modeled CME - evolve with increasing heliocentric distance.** 3-D MHD modeling of the solar wind structures associated with the 13 December 2006 coronal mass ejection event (Fig. 1).




Fig 1: Kataoka et al. 2009


■ 3D structure of ICME and solar wind

- **ICMEs (Interplanetary Coronal Mass Ejections) - sweep out interplanetary space plasmas and magnetic fields.** ICMEs can coalesce with other ICMEs and high speed streams/CIRs.
- **Multiple flaring and multiple CME releases** → a very **complex solar wind** → **deflection and retardation of ICME propagation.**
- **Development of improved mean field solar dynamo models** taking into account multi-cellular and/or time varying meridional circulation, magnetic pumping, North-South asymmetry, irregularity in the cycles.

■ Other Scientific Highlights

- **Numerical Investigation of a Coronal Mass Ejection** from an Anemone Active Region: Reconnection and Deflection of the 2005 August 22 Eruption
- **Magneto-thermal convection in solar prominences** observed by *Hinode* satellite.
- **Superflares on solar type stars** - discovered by *Kepler* mission.

E-SCIENCE AND INFORMATICS



- Worldwide attention to data-aspects of international collaborations
 - good
- Relations to the World Data System
 - okay
- Use of virtual observatories as well as localized data sharing
 - very good

CAWSES


http://www.cawses.org/wiki/index.php/Main_Page

CAWSES

page discussion edit history delete move protect unwatch

Main Page

Climate And Weather of the Sun-Earth System (CAWSES 2009-2013) is an international program sponsored by SCOSTEP (Scientific Committee on Solar-Terrestrial Physics) established with an aim of significantly enhancing our understanding of the space environment and its impacts on life and society. The main functions of CAWSES are to help coordinate international activities in observations, modeling, and applications crucial to achieving this understanding, to involve scientists in both developed and developing countries, and to provide educational opportunities for students of all levels.



navigation

- Main Page
- Task Groups
- Conferences
- Capacity Building
- Virtual Institute
- Publications
- News
- Steering Committee
- Help

search

Go Search

toolbox

- What links here
- Related changes
- Upload file
- Special pages
- Printable version
- Permanent link

This page was last modified on 1 September 2009, at 15:04. This page has been accessed 1,750 times. Privacy policy About CAWSES

CAWSES Wiki

http://www.cawses.org/wiki/index.php/Main_Page

Task Groups – CAWSES

http://www.cawses.org/wiki/index.php/Task_Groups

Task Groups – CAWSES

page discussion edit history delete move protect unwatch

Task Groups

CAWSES II is organized around Task Groups

Contents [hide]

- 1 Task 1: What are the solar influences on the Earth's climate?
- 2 Task 2: How will geospace respond to an altered climate?
- 3 Task 3: How does short-term solar variability affect the geospace environment?
- 4 Task 4: What is the geospace response to variable waves from the lower atmosphere?
- 5 Capacity Building
- 6 Virtual Institute

Task 1: What are the solar influences on the Earth's climate?

Solar variability drives geo-space and the atmosphere on time scales ranging from minutes to millennia. Feedbacks are inherent in the Earth system and may amplify the direct forcing effects from the Sun. The influence of this solar variability on Earth's climate is a key issue of the Intergovernmental Panel on Climate Change, and one that continues to be highlighted by policy makers, climate change skeptics, and the media.

Task 2: How will geospace respond to an altered climate?

Radiative, chemical, and dynamical forcing from below contributes to disturbances of the upper atmosphere. In response to rising greenhouse gas concentrations, cooling in the middle atmosphere will alter the complex physical and chemical processes of this region. Patterns of cooling and contraction of the upper atmosphere are emerging from model studies and observations, consistent with a strong connection to changes in the lower atmosphere. Recent changes in noctilucent cloud distributions, now observed on a global scale by the AIM satellite (see figure above), are thought to be symptomatic of cooling temperatures in the upper atmosphere. Rising greenhouse gas concentrations alter the ionosphere in a variety of ways and could be transmitted to the magnetosphere. These changes may have unforeseen consequences for space-related technologies and societal infrastructures.

Task 3: How does short-term solar variability affect the geospace environment?

Solar variations such as solar flares, energetic particle bursts, coronal mass ejections, and high-speed solar wind streams directly affect the geospace environment. Electromagnetic disturbances in the ionosphere, while not directly related to climate, can affect the

navigation

- Main Page
- Task Groups
- Conferences
- Capacity Building
- Virtual Institute
- Publications
- News
- Steering Committee
- Help

search

Go Search

toolbox

- What links here
- Related changes
- Upload file
- Special pages
- Printable version
- Permanent link

Wiki in place supporting science and outreach tasks – a place to collaborate

- 2 of 4 task groups made extensive use of the wiki
- Other 2 groups documented what they were doing



Thank You!